


# Fifth Metacarpal Neck Fracture (Boxer's Fracture)

## What is a fractured metacarpal?

The metacarpals are bones in the hand which extend from the knuckles to the wrist. Some of the most common metacarpal fractures (breaks) involve the fifth (small) bone the hand and occurs when the bone just below the knuckle of the little finger is broken.

It is also called a 'boxer's fracture' and is usually caused by the impact of a clenched fist with a hard immovable object. The knuckle compresses and it snaps the neck of the metacarpal bone. The break will heal, and the pain will usually settle in three weeks.



## What are the symptoms?

Patients with metacarpal fractures experience pain, swelling, loss of gripping power and difficulty in moving the fingers, especially over the affected metacarpal bones. The knuckle may look as if it has "disappeared" due to the focal swelling and angulation of the fracture.

## What is the treatment?

- The fracture usually heals by itself, so the treatment is aimed at making it a bit more comfortable. For most people this involves strapping the little finger to the ring finger (neighbour strapping) and a Velcro wrist splint which can be worn for up to 14 days. The finger strapping may help the pain and allow early movement.
- The fracture site can be painful and swollen, especially for the first week. Therefore, it is important for you to keep it elevated. Sometimes you will be given a sling to wear to help reduce the swelling.

continued over...

- 
- 
- There is a risk that heavy lifting may be sore for 6 or 8 weeks.
  - Taking regular simple pain killers will be helpful in relieving your pain.
  - Smoking is known to delay healing and can prevent it all together. If possible, we would advise you to stop. Further advice can be found on the NHS website <https://www.nhs.uk/better-health/quit-smoking/> or from your GP
  - You will be referred to the Hand Clinic after your initial management at the Minor Injury Unit.

### **Is there anything I can do to help?**

- It is important in the first few days that you try and get your fingers moving as much as possible. If your fingers become stiff, they will be very difficult to get moving again. Please try to make a full fist as soon as the pain allows and repeat this regularly, during the course of your healing and recovery.
- Use your hand as normally as possible within the limits of the pain. This will not cause further damage, but failure to do so may lead to ongoing stiffness.

### **How long will it take before I am able to use my hand as usual?**


The fracture itself does not heal for about 6 weeks, so heavier work and contact sport must be left until then.

### **What should I expect in the long term?**

- Handgrip is generally very good after this type of injury.
- There will be a lump on the back of the hand near the knuckle of the little finger. This is normal and occurs partly because the broken bone has bent slightly. It also develops as the body forms new bone around the fracture to help it heal. This lump should smooth out over a year or two and will not cause any problems with the use of your hand.
- There will be some shortening expected as a consequence of the injury. The finger may droop a little, but this will usually recover over time.


### **Is there anything I should look out for?**

It is normal to experience some pain as well as some stiffness and a weakened grip, but these symptoms should gradually improve. However, if the pain gets worse or lasts more than 3 weeks or you are concerned about your hand, please contact the Fracture clinic - 0151 604 7069 (Monday to Friday, 8.30am until 3.30 pm)



---

If your little finger does not tuck in neatly next to the ring finger when making a fist, there is a risk that it may be malrotated. If so, please contact the Fracture Clinic.



**Please contact your GP or NHS111 for any queries or concerns.**

**Please retain this leaflet for future reference.**

**Reference:**

Fifth Metacarpal Neck Fracture (Boxer's Fracture) Leaflet, Orthopaedic Department, Wirral University Teaching Hospital NHS Foundation Trust

If you would like this information in another format or language, please contact the Your Experience Team on freephone 0800 694 5530. Alternatively you can email [wcnt.yourexperience@nhs.net](mailto:wcnt.yourexperience@nhs.net)

September 2021 V2